

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
4 October 2001 (04.10.2001)

PCT

(10) International Publication Number
WO 01/72714 A2

(51) International Patent Classification⁷: **C07D 231/00**

(21) International Application Number: **PCT/US01/08972**

(22) International Filing Date: **22 March 2001 (22.03.2001)**

(25) Filing Language: **English**

(26) Publication Language: **English**

(30) Priority Data:
60/191,757 **24 March 2000 (24.03.2000)** **US**

(71) Applicant (for all designated States except US): **COCEN-SYS, INC.** [US/US]; 213 Technology Drive, Irvine, CA 92618 (US).

(72) Inventors; and
(75) Inventors/Applicants (for US only): **HOGENKAMP, Derk** [US/US]; 4510 Park Drive, Carlsbad, CA 92008 (US). **NGUYEN, Phong** [US/US]; 1048 Wingfoot Street, Placentia, CA 92870 (US). **YANG, Ji** [CN/US]; 2010 Quail Ridge Drive, Plainsboro, NJ 08536 (US).

(74) Agents: **COVERT, John, M.** et al.; Sterne, Kessler, Goldstein & Fox P.L.L.C., 1100 New York Avenue, N.W., Suite 600, Washington, DC 20005-3934 (US).

(81) Designated States (national): **AE, AG, AI, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.**

(84) Designated States (regional): **ARIPO** patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), **Eurasian** patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), **European** patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), **OAPI** patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published:


— without international search report and to be republished upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.



WO 01/72714 A2

(54) Title: **ARYL SUBSTITUTED PYRAZOLES, TRIAZOLES AND TETRAZOLES, AND THE USE THEREOF**



(57) Abstract: This invention relates to compounds having the Formula (I) or a pharmaceutically acceptable salt, prodrug or solvate thereof, wherein Het and R₅ - R₈ are set in the specification. The invention also is directed to the use of compounds of Formula (I) for the treatment of neuronal damage following global and focal ischemia, for the treatment or prevention of neurodegenerative conditions such as amyotrophic lateral sclerosis (ALS), and for the treatment, prevention or amelioration of both acute or chronic pain, as antitinnitus agents, as anticonvulsants, and as antimanic depressants, as local anesthetics, as antiarrhythmics and for the treatment or prevention of diabetic neuropathy.

21567
#2

Aryl Substituted Pyrazoles, Triazoles and Tetrazoles, and the Use Thereof

Background of the Invention

5

Field of the Invention

This invention is in the field of medicinal chemistry. In particular, the invention relates to aryl substituted pyrazoles, triazoles and tetrazoles, and the discovery that these compounds are anticonvulsants and act as blockers of sodium (Na⁺) channels.

Related Background Art

Several classes of therapeutically useful drugs, including local anesthetics such as lidocaine and bupivacaine, antiarrhythmics such as propafenone and amiodarone, and anticonvulsants such as lamotrigine, phenytoin and carbamazepine, have been shown to share a common mechanism of action by blocking or modulating Na⁺ channel activity (Catterall, W.A., *Trends Pharmacol. Sci.* 8:57-65 (1987)). Each of these agents is believed to act by interfering with the rapid influx of Na⁺ ions.

Recently, other Na⁺ channel blockers such as BW619C89 and lifarizine have been shown to be neuroprotective in animal models of global and focal ischemia and are presently in clinical trials (Graham *et al.*, *J. Pharmacol. Exp. Ther.* 269:854-859 (1994); Brown *et al.*, *British J. Pharmacol.* 115:1425-1432 (1995)).


The neuroprotective activity of Na⁺ channel blockers is due to their effectiveness in decreasing extracellular glutamate concentration during ischemia by inhibiting the release of this excitotoxic amino acid neurotransmitter. Studies have shown that unlike glutamate receptor antagonists, Na⁺ channel blockers prevent hypoxic damage to mammalian white matter (Stys *et al.*, *J. Neurosci.* 12:430-439 (1992)). Thus, they may offer advantages for treating certain types of strokes or neuronal trauma where damage to white matter tracts is prominent.

- 2 -

Another example of clinical use of a Na⁺ channel blocker is riluzole. This drug has been shown to prolong survival in a subset of patients with ALS (Bensim *et al.*, *New Engl. J. Med.* 330:585-591 (1994)) and has subsequently been approved by the FDA for the treatment of ALS. In addition to the above-mentioned clinical uses, carbamazepine, lidocaine and phenytoin are occasionally used to treat neuropathic pain, such as from trigeminal neurologia, diabetic neuropathy and other forms of nerve damage (Taylor and Meldrum, *Trends Pharmacol. Sci.* 16:309-316 (1995)), and carbamazepine and lamotrigine have been used for the treatment of manic depression (Denicott *et al.*, *J. Clin. Psychiatry* 55: 70-76 (1994)). Furthermore, based on a number of similarities between chronic pain and tinnitus (Moller, A. R. *Am. J. Otol.* 18: 577-585 (1997); Tonndorf, *J. Hear. Res.* 28: 271-275 (1987)) it has been proposed that tinnitus should be viewed as a form of chronic pain sensation (Simpson, J. J. and Davies, E. W. *Tip.* 20: 12-18 (1999)). Indeed, lignocaine and carbamazepine have been shown to be efficacious in treating tinnitus (Majumdar, B. *et al. Clin. Otolaryngol.* 8: 175-180 (1983); Donaldson, I. *Laryngol. Otol.* 95: 947-951 (1981)).

It has been established that there are at least five to six sites on the voltage-sensitive Na⁺ channels which bind neurotoxins specifically (Catterall, W.A., *Science* 242:50-61 (1988)). Studies have further revealed that therapeutic antiarrhythmics, anticonvulsants and local anesthetics whose actions are mediated by Na⁺ channels, exert their action by interacting with the intracellular side of the Na⁺ channel and allosterically inhibiting interaction with neurotoxin receptor site 2 (Catterall, W.A., *Ann. Rev. Pharmacol. Toxicol.* 10:15-43 (1980)).

Cocco, M. T.; Maccioni, A.; Plumitallo, A.; *Farmaco Ed.Sci.*; 40: 1985; 272-284 describe the two following compounds:



Other aryl substituted heterocycles are described in Stefancich, G. *et al. Arch.Pharm.(Weinheim Ger.)* 323: 273-280 (1990) as antimycotic agents. These compounds are described in Appendix A.

Compounds of Formula *I* have not been used heretofor for treating a disorder responsive to the blockade of sodium channels in a mammal.

Summary of the Invention

The present invention is related to the discovery that aryl substituted pyrazoles, triazoles and tetrazoles represented by Formula *I* act as blockers of sodium (Na^+) channels.

The invention is also related with treating a disorder responsive to the blockade of sodium channels in a mammal suffering from excess activity of said channels by administering an effective amount of a compound of Formula *I* as described herein.

A further aspect of the present invention is to provide a method for treating, preventing or ameliorating neuronal loss following global and focal ischemia; treating, preventing or ameliorating pain including acute and chronic pain, and neuropathic pain; treating, preventing or ameliorating convulsion and neurodegenerative conditions; treating, preventing or ameliorating manic depression; using as local anesthetics, antiarrhythmics, and treating tinnitus by administering a compound of Formula *I* to a mammal in need of such treatment.

Another aspect of the present invention is directed to the use of the compounds of Formula *I* as blockers of sodium channels.

The present invention is also directed to the use of a compound of Formula *I* for the treatment of neuronal damage following global and focal ischemia, and for the treatment or prevention of neurodegenerative conditions, such as amyotrophic lateral sclerosis (ALS), for the treatment of tinnitus, as antimanic depressants, as local anesthetics, as antiarrhythmics, as anticonvulsants and for the treatment or prevention

- 4 -

of diabetic neuropathy and for the treatment of pain including both acute and chronic pain and migraine headache.

5 A further aspect of the present invention is to provide a pharmaceutical composition useful for treating disorders responsive to the blockade of sodium ion channels, containing an effective amount of a compound of Formula *I* in a mixture with one or more pharmaceutically acceptable carriers or diluents.

A number of compounds useful in the present invention have not been heretofor reported. Thus, the present invention is also directed to novel aryl substituted pyrazoles, triazoles and tetrazoles of Formula *I*.

10 Further, the present invention is directed to ^3H and ^{14}C radiolabeled compounds of Formula *I* and their use as radioligands for their binding site on the sodium channel.

15 Additional embodiments and advantages of the invention will be set forth in the description which follows, and in part will be obvious from the description, or may be learned by practice of the invention. The embodiments and advantages of the invention will be realized and attained by means of the elements and combinations particularly pointed out in the appended claims.

20 It is to be understood that both the foregoing general description and the following detailed description are exemplary and explanatory only and are not restrictive of the invention, as claimed.


Detailed Description of the Invention

25 The present invention arises out of the discovery that the aryl substituted pyrazoles, triazoles and tetrazoles of Formula *I* act as blockers of Na^+ channels. In view of this discovery, compounds of Formula *I* are useful for treating disorders responsive to the blockade of sodium ion channels.

The compounds useful in this aspect of the present invention are the aryl substituted pyrazoles, triazoles and tetrazoles represented by Formula *I*:

30

- 5 -




I

or a pharmaceutically acceptable salt, prodrug or solvate thereof, wherein

X is one of O, S, NR₉, CH₂, NR₉C(O), or C(O)NR₉, where R₉ is hydrogen or C₁-C₁₀ alkyl;

5 Het is a heteroaryl selected from the group consisting of



10 R₁ is selected from the group consisting of hydrogen, optionally substituted alkyl, optionally substituted heteroaryl, C(O)R₁₀, CH₂C(O)R₁₀, S(O)R₁₀, and SO₂R₁₀;

R₂ and R₃ are independently selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, aryl, cyano, aminoalkyl, hydroxyalkyl, alkoxyalkyl, alkylthio, alkylsulfinyl, alkylsulfonyl, carboxyalkyl, alkylamino, dialkylamino, aminocarbonyl, alkylaminocarbonyl, arylaminocarbonyl, aralkylaminocarbonyl, alkylcarbonylamino, arylcarbonylamino, aralkylcarbonylamino, alkylcarbonyl, aminosulfonyl, alkylaminosulfonyl, and alkylsulfonyl;

20 R₅, R₆, R₇, and R₈ are independently selected from the group consisting of hydrogen, halo, haloalkyl, alkyl, alkenyl, alkynyl, hydroxyalkyl, aminoalkyl, carboxyalkyl, alkoxyalkyl, nitro, amino, ureido, cyano, acylamino, amide, hydroxy, thiol, acyloxy, azido, alkoxy, carboxy, carbonylamido and alkylthiol;

R₁₀ is selected from the group consisting of amino, alkyl, alkenyl, alkynyl, OR₁₁, alkylamino, dialkylamino, alkenylamino, dialkylaminoalkenyl, cycloalkyl, aryl, heterocycle, heteroaryl, aralkyl, arylalkenyl, arylalkynyl, and cycloalkylalkylamino;

- 6 -

R_{11} is selected from the group consisting of hydrogen, optionally substituted alkyl, and an alkalimetal.

Thus, the present invention is directed to provide a method for treating, preventing or ameliorating neuronal loss following global and focal ischemia; treating, preventing or ameliorating pain including acute and chronic pain, and neuropathic pain; treating, preventing or ameliorating convulsion and neurodegenerative conditions; treating, preventing or ameliorating manic depression; using as local anesthetics, antiarrhythmics, and treating tinnitus by administering a compound of Formula I to a mammal in need of such treatment.

The present invention is also directed to novel compounds having Formula I as described above; provided that:

- 1) when Het is (ii), and X is O, then R_{10} is not alkyl, aralkyl, aryl, or OR_{11} ;
- 2) when Het is (i) or (ii), then X is not NR_9 ;
- 3) when Het is (iii), then X is not CH_2 ; and
- 4) when Het is (iii), and X is O, then R_{10} is not OR_{11} .

Optional substituents on optionally substituted groups, when not otherwise defined, include one or more groups independently selected from the group consisting of halo, halo(C_{1-6}) alkyl, aryl, pyrimidine, cycloalkyl, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, aryl(C_{1-6})alkyl, aryl(C_{2-6})alkenyl, aryl(C_{2-6})alkynyl, cycloalkyl(C_{1-6})alkyl, hydroxy(C_{1-6})alkyl, amino(C_{1-6})alkyl, carboxy(C_{1-6})alkyl, alkoxy(C_{1-6})alkyl, nitro, amino, ureido, cyano, C_{1-6} acylamino, hydroxy, thiol, C_{1-6} acyloxy, azido, C_{1-6} alkoxy, carboxy, aminocarbonyl, carbamoyloxy, C_{1-6} alkylsulfonylamino, C_{1-6} acyl, and C_{1-6} alkylthiol groups mentioned above as long as the resulting compound is stable. Preferred optional substituents include: halo, halo(C_{1-6})alkyl, hydroxy(C_{1-6})alkyl, amino(C_{1-6})alkyl, hydroxy, nitro, C_{1-6} alkyl, C_{1-6} alkoxy, aminocarbonyl, carbamoyloxy, C_{1-6} alkylsulfonylamino, C_{1-6} acyl and amino.

Preferably, R_1 is selected from the group consisting of an alkyl optionally substituted by halogen, hydroxy, carbamoyloxy, C_{1-6} acyl, C_{1-6} alkylsulfonylamino, aryl, preferably phenyl, or aminocarbonyl; $C(O)R_{10}$; $CH_2C(O)R_{10}$; or SO_2R_{10} , wherein R_{10} is selected from the group consisting of C_{1-6} alkyl, C_{2-6} alkenyl, OR_{11} , amino, C_{1-6}

- 7 -

alkylamino, di(C₁₋₆)alkylamino, C₂₋₆ alkenylamino, heterocycle and mono- and di-(C₁₋₆)alkylaminoalkenyl, and wherein R₁₁ is as defined above.

Preferably, R₁₀ is selected from the group consisting of C₁₋₆ alkyl, C₂₋₆ alkenyl, OR₁₁, amino, C₁₋₆ alkylamino, di(C₁₋₆)alkylamino, C₂₋₆ alkenylamino, mono- and di-(C₁₋₆)alkylamino(C₂₋₆)alkenyl, N-morpholinyl, N-pyrrolidinyl, and N-piperazinyl, all of which can be optionally substituted, wherein R₁₁ is as defined above.

Preferably, R₂ and R₃ are independently selected from the group consisting of hydrogen, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, amino(C₁₋₆)alkyl, amino, cyano, C₁₋₆ alkoxy, C₁₋₆ alkylthio, C₁₋₆ alkylsulfinyl, hydroxy(C₁₋₆)alkyl, alkoxy(C₁₋₆)alkyl, carboxy(C₁₋₆)alkyl, aminocarbonyl, C₁₋₆ alkylaminocarbonyl, C₆₋₁₀ arylaminocarbonyl, C₆₋₁₀ aryl(C₁₋₆)alkylaminocarbonyl, C₁₋₆ alkylcarbonylamino, C₆₋₁₀ arylcarbonylamino, and C₆₋₁₀ aryl(C₁₋₆)alkylcarbonylamino, more preferably hydrogen, C₁₋₆ alkyl, C₁₋₆ alkoxy, amino(C₁₋₆)alkyl, C₁₋₆ alkylthio and aminocarbonyl.

The groups R₅-R₈ each take place of a hydrogen atom that would otherwise be present in any position on the aryl ring to which the R group is attached.

Preferably, R₅, R₆, R₇, and R₈ are independently selected from the group consisting of hydrogen, halo (preferably chloro or fluoro), halo(C₁₋₆)alkyl, C₁₋₆ alkyl, hydroxy(C₁₋₆)alkyl, amino(C₁₋₆)alkyl, carboxy(C₁₋₆)alkyl, alkoxy(C₁₋₆)alkyl, nitro, amino, C₁₋₆ acylamino, amide, hydroxy, thiol, C₁₋₆ acyloxy, C₁₋₆ alkoxy, carboxy, carbonylamido and C₁₋₆ alkylthiol.

One group of preferred compounds falling within the scope of Formula I include compounds wherein R₁ is C(O)R₁₀ or SO₂R₁₀, where R₁₀ is defined above, and is more preferably amino or C₁₋₆ alkyl. In this group of compounds, X is more preferably O or S, most preferably O.

Especially preferred in this group are compounds where R₅ and R₆ are each hydrogen; R₂ and R₃ are both H; and R₇ and R₈ are selected from the group consisting of hydrogen, halo, halo(C₁₋₆)alkyl, C₁₋₆ alkyl, hydroxy(C₁₋₆)alkyl, amino(C₁₋₆)alkyl, carboxy(C₁₋₆)alkyl, alkoxy(C₁₋₆)alkyl, nitro, amino, C₁₋₆

- 8 -

acylamino, amide, hydroxy, thiol, C₁-C₆ acyloxy, C₁-C₆ alkoxy, carboxy, carbonylamido and C₁-C₆ alkylthiol.

Another group of preferred compounds includes compounds of Formula I:




5

or a pharmaceutically acceptable salt, prodrug or solvate thereof, wherein

X is O or S, preferably O;

Het is a heteroaryl selected from the group consisting of



10

preferably (i) or (iii);

R₁ is C(O)R₁₀, CH₂C(O)R₁₀, or SO₂R₁₀ wherein R₁₀ is amino, alkyl, N-morpholinyl, N-pyrrolidinyl or N-piperazinyl, more preferably amino, all of which can be optionally substituted. R₁ is preferably C(O)R₁₀, where R₁₀ is amino, C₁-C₆ alkyl, or a heterocycle, such as N-morpholinyl, N-pyrrolidinyl and N-piperazinyl;

15

R₂ and R₃ are independently hydrogen, C₁-C₆ alkyl, C₁-C₆ alkylthio or C₁-C₆ alkylsulfinyl, with R₂ and R₃ preferably being hydrogen; R₅ and R₆ are as defined above and are preferably hydrogen; and R₇ and R₈ are independently selected from the group consisting of hydrogen, halo, halo(C₁-C₆)alkyl, C₁-C₆ alkyl, hydroxy(C₁-C₆)alkyl, amino(C₁-C₆)alkyl, carboxy(C₁-C₆)alkyl, alkoxy(C₁-C₆)alkyl, nitro, amino, C₁-C₆ acylamino, amide, hydroxy, thiol, C₁-C₆ acyloxy, C₁-C₆ alkoxy, carboxy, carbonylamido and C₁-C₆ alkylthiol;

20

provided that:

- 1) when Het is (ii), and X is O, then R₁₀ is not alkyl, aralkyl, aryl or OR₁₁;
- 2) when Het is (iii), and X is O, then R₁₀ is not OR₁₁.

Exemplary preferred compounds that may be employed in this method of invention include, without limitation:

1-[4-(4-nitrophenoxy)phenyl]-1H-[1,2,4]triazole;
1-[4-(4-fluorophenoxy)phenyl]-3-methylpyrazole;
3-methyl-1-(4-phenoxyphenyl)pyrazole;
1-(4-phenoxyphenyl)-1H-pyrazole-3-carboxamide;
1-(4-phenoxyphenyl)-1H-pyrazole-5-carboxamide;
1-[4-(4-fluorophenoxy)phenyl]-1H-pyrazole-3-carboxamide;
1-[4-(4-nitrophenoxy)phenyl]-1H-[1,2,4]triazole-3-carboxamide; and
1-[4-(4-chloro-2-fluorophenoxy)phenyl]-1H-pyrazole-3-carboxamide.

Another group of exemplary preferred compounds that may be employed in this invention include 1-[4-(4-fluorophenoxy)phenyl]-5-methylpyrazole, 1-(4-phenoxyphenyl)-1H-pyrazole-4-carboxamide, and 4-(4-fluorophenoxy)phenylpyrazole.

Useful aryl groups are C₆₋₁₄ aryl, especially C₆₋₁₀ aryl. Typical C₆₋₁₄ aryl groups include phenyl, naphthyl, phenanthryl, anthracyl, indenyl, azulenyl, biphenyl, biphenylenyl and fluorenyl groups.

Useful cycloalkyl groups are C₃₋₈ cycloalkyl. Typical cycloalkyl groups include cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl and cycloheptyl.

The term "heteroaryl" as employed herein refers to groups having 5 to 14 ring atoms; 6, 10 or 14 π electrons shared in a cyclic array; and containing carbon atoms and 1, 2 or 3 oxygen, nitrogen or sulfur heteroatoms (where examples of heteroaryl groups are: thienyl, benzo[b]thienyl, naphtho[2,3-b]thienyl, thianthrenyl, furyl, benzofuryl, pyranyl, isobenzofuranyl, benzoxazolyl, chromenyl, xanthenyl, phenoxathiinyl, 2H-pyrrolyl, pyrrolyl, imidazolyl, pyrazolyl, pyridyl, pyrazinyl, pyrimidinyl, pyridazinyl, indoliziny, isoindolyl, 3H-indolyl, indolyl, indazolyl, purinyl, 4H-quinoliziny, isoquinolyl, quinolyl, phthalazinyl, naphthyridinyl, quinazolinyl, cinnolinyl, pteridinyl, 4aH-carbazolyl, carbazolyl, β -carbolinyl,

phenanthridinyl, acridinyl, perimidinyl, phenanthrolinyl, phenazinyl, thiazolyl, isothiazolyl, phenothiazinyl, isoxazolyl, furazanyl, and phenoxazinyl groups).

Useful halo or halogen groups include fluorine, chlorine, bromine and iodine.

5 Useful alkyl groups include straight-chained and branched C₁₋₁₀ alkyl groups, more preferably C₁₋₆ alkyl groups. Typical C₁₋₁₀ alkyl groups include methyl, ethyl, propyl, isopropyl, butyl, *sec*-butyl, *tert*-butyl, 3-pentyl, hexyl and octyl groups. Also contemplated is a trimethylene group substituted on two adjoining positions on the benzene ring of the compounds of the invention.

10 Useful alkenyl groups are C₂₋₆ alkenyl groups, preferably C₂₋₄ alkenyl. Typical C₂₋₄ alkenyl groups include ethenyl, propenyl, isopropenyl, butenyl, and *sec*-butenyl.

Useful alkynyl groups are C₂₋₆ alkynyl groups, preferably C₂₋₄ alkynyl. Typical C₂₋₄ alkynyl groups include ethynyl, propynyl, butynyl, and 2-butynyl groups.

15 Useful arylalkyl groups include any of the above-mentioned C₁₋₁₀ alkyl groups substituted by any of the above-mentioned C₆₋₁₄ aryl groups. Useful values include benzyl, phenethyl and naphthylmethyl.

Useful arylalkenyl groups include any of the above-mentioned C₂₋₄ alkenyl groups substituted by any of the above-mentioned C₆₋₁₄ aryl groups.

20 Useful arylalkynyl groups include any of the above-mentioned C₂₋₄ alkynyl groups substituted by any of the above-mentioned C₆₋₁₄ aryl groups. Useful values include phenylethynyl and phenylpropynyl.

Useful heteroarylalkyl groups include any of the above-mentioned C₁₋₁₀ alkyl groups substituted by any of the above-mentioned heteroaryl groups.

Useful heteroarylalkenyl groups include any of the above-mentioned C₂₋₄ alkenyl groups substituted by any of the above-mentioned heteroaryl groups.

25 Useful heteroarylalkynyl groups include any of the above-mentioned C₂₋₄ alkynyl groups substituted by any of the above-mentioned heteroaryl groups.

Useful cycloalkylalkyl groups include any of the above-mentioned C₁₋₁₀ alkyl groups substituted by any of the above-mentioned cycloalkyl groups.

Useful haloalkyl groups include C_{1-10} alkyl groups substituted by one or more fluorine, chlorine, bromine or iodine atoms, e.g. fluoromethyl, difluoromethyl, trifluoromethyl, pentafluoroethyl, 1,1-difluoroethyl and trichloromethyl groups.

5 Useful hydroxyalkyl groups include C_{1-10} alkyl groups substituted by hydroxy, e.g. hydroxymethyl, hydroxyethyl, hydroxypropyl and hydroxybutyl groups.

Useful alkoxy groups include oxygen substituted by one of the C_{1-10} alkyl groups mentioned above.

Useful alkylthio groups include sulfur substituted by one of the C_{1-10} alkyl groups mentioned above.

10 Useful acylamino groups are any C_{1-6} acyl (alkanoyl) attached to an amino nitrogen, e.g. acetamido, propionamido, butanoylamido, pentanoylamido, hexanoylamido as well as aryl-substituted C_{2-6} substituted acyl groups.

Useful acyloxy groups are any C_{1-6} acyl (alkanoyl) attached to an oxy (-O-) group, e.g. acetoxy, propionyloxy, butanoyloxy, pentanoyloxy, hexanoyloxy and the like.

15 The term heterocycle is used herein to mean saturated or partially unsaturated 3-7 membered monocyclic, or 7-10 membered bicyclic ring system, which consists of carbon atoms and from one to four heteroatoms independently selected from the group consisting of O, N, and S, wherein the nitrogen and sulfur heteroatoms can be optionally oxidized, the nitrogen can be optionally quaternized, and including any bicyclic group in which any of the above-defined heterocyclic rings is fused to a benzene ring, and wherein the heterocyclic ring can be substituted on carbon or on a nitrogen atom if the resulting compound is stable. Examples include, but are not limited to, pyrrolidine, piperazine, morpholine, imidazoline, pyrazolidine, benzodiazepines and the like.

25 Useful heterocycloalkyl groups include any of the above-mentioned C_{1-10} alkyl groups substituted by any of the above-mentioned heterocyclic groups.

Useful alkylamino and dialkylamino groups are —NHR_{20} and $\text{—NR}_{20}\text{R}_{21}$, wherein R_{20} and R_{21} are C_{1-10} alkyl groups.

Aminocarbonyl group is —C(O)NH_2 .

Useful alkylaminocarbonyl groups are carbonyl groups substituted by —NHR_{20} and $\text{—NR}_{20}\text{R}_{21}$, wherein R_{20} and R_{21} are C_{1-10} alkyl groups as defined above.

Useful alkylthiol groups include any of the above-mentioned C_{1-10} alkyl groups substituted by a —SH group.

Useful alkylsulfinyl groups include any of the above-mentioned C_{1-10} alkyl groups attached to a sulfinyl (—SO—).

Useful alkylsulfonyl groups include any of the above-mentioned C_{1-10} alkyl groups attached to a sulfonyl ($\text{—SO}_2\text{—}$).

A carbamoyloxy group is —O—C(O)—NH_2 .

A carboxy group is —COOH .

An azido group is —N_3 .

An ureido group is —NH—C(O)—NH_2 .

An amino group is —NH_2 .

An amide group is an organic radical having —NHC(O)— as a functional group.

The invention disclosed herein is meant to encompass all pharmaceutically acceptable salts thereof of the disclosed compounds. The pharmaceutically acceptable salts include, but are not limited to, metal salts such as sodium salt, potassium salt, cesium salt and the like; alkaline earth metals such as calcium salt, magnesium salt and the like; organic amine salts such as triethylamine salt, pyridine salt, picoline salt, ethanolamine salt, triethanolamine salt, dicyclohexylamine salt, N,N' -dibenzylethylenediamine salt and the like; inorganic acid salts such as hydrochloride, hydrobromide, sulfate, phosphate and the like; organic acid salts such as formate, acetate, trifluoroacetate, maleate, tartrate and the like; sulfonates such as methanesulfonate, benzenesulfonate, p-toluenesulfonate, and the like; amino acid salts such as arginate, aspartate, glutamate and the like.

The invention disclosed herein is also meant to encompass prodrugs of the disclosed compounds. Prodrugs are considered to be any covalently bonded carriers which release the active parent drug *in vivo*. Examples of prodrugs include esters or

amides of Formula I with R_2-R_{11} as hydroxyalkyl or aminoalkyl, and these may be prepared by reacting such compounds with anhydrides such as succinic anhydride.

The invention disclosed herein is also meant to encompass the *in vivo* metabolic products of the disclosed compounds. Such products may result for example from the oxidation, reduction, hydrolysis, amidation, esterification and the like of the administered compound, primarily due to enzymatic processes. Accordingly, the invention includes compounds produced by a process comprising contacting a compound of this invention with a mammal for a period of time sufficient to yield a metabolic product thereof. Such products typically are identified by preparing a radiolabelled compound of the invention, administering it parenterally in a detectable dose to an animal such as rat, mouse, guinea pig, monkey, or to man, allowing sufficient time for metabolism to occur and isolating its conversion products from the urine, blood or other biological samples.

The invention disclosed herein is also meant to encompass the disclosed compounds being isotopically-labelled by having one or more atoms replaced by an atom having a different atomic mass or mass number. Examples of isotopes that can be incorporated into the disclosed compounds include isotopes of hydrogen, carbon, nitrogen, oxygen, phosphorous, fluorine and chlorine, such as ^2H , ^3H , ^{13}C , ^{14}C , ^{15}N , ^{18}O , ^{17}O , ^{31}P , ^{32}P , ^{35}S , ^{18}F , and ^{36}Cl , respectively.

Some of the compounds disclosed herein may contain one or more asymmetric centers and may thus give rise to enantiomers, diastereomers, and other stereoisomeric forms. The present invention is also meant to encompass racemic mixtures, resolved forms mixtures thereof, as well as the individual enantiomers that may be separated according to methods that are well known to those of ordinary skill in the art. When the compounds described herein contain olefinic double bonds or other centers of geometric asymmetry, and unless specified otherwise, it is intended to include both E and Z geometric isomers. All tautomers are intended to be encompassed by the present invention as well.

As used herein, the term "stereoisomers" is a general term for all isomers of individual molecules that differ only in the orientation of their atoms in space. It

- 14 -

includes enantiomers and isomers of compounds with more than one chiral center that are not mirror images of one another (diastereomers).

The term "chiral center" refers to a carbon atom to which four different groups are attached.

5 The term "enantiomer" or "enantiomeric" refers to a molecule that is nonsuperimposeable on its mirror image and hence optically active wherein the enantiomer rotates the plane of polarized light in one direction and its mirror image rotates the plane of polarized light in the opposite direction.

10 The term "racemic" refers to a mixture of equal parts of enantiomers and which is optically inactive.

The term "resolution" refers to the separation or concentration or depletion of one of the two enantiomeric forms of a molecule. The phrase "enantiomeric excess" refers to a mixture wherein one enantiomer is present in a greater concentration than its mirror image molecule.

15 Since the compounds of Formula *I* are blockers of sodium (Na^+) channels, a number of diseases and conditions mediated by sodium ion influx can be treated employing these compounds. Therefore, the invention is related to a method of treating, preventing or ameliorating neuronal loss associated with stroke, global and focal ischemia, CNS trauma, hypoglycemia and surgery, spinal cord trauma; as well
20 as treating or ameliorating neurodegenerative diseases including Alzheimer's disease, amyotrophic lateral sclerosis, Parkinson's disease, treating or ameliorating anxiety, convulsions, glaucoma, migraine headache, and muscle spasm. The compounds of Formula *I* are also useful as antitinnitus agents, antimanic depressants, as local anesthetics, and as antiarrhythmics; as well as for treating, preventing or ameliorating
25 pain including surgical, chronic and neuropathic pain. In each instance, the methods of the present invention require administering to an animal in need of such treatment an effective amount of a sodium channel blocker of the present invention, or a pharmaceutically acceptable salt or prodrug thereof.


30 The invention is also directed to a method for treating disorders responsive to the blockade of sodium channels in animals suffering thereof. Particular preferred

embodiments of the aryl substituted heteroaryl compounds for use in method of this invention are represented by previously defined Formula I.

The compounds of this invention may be prepared using methods known to those skilled in the art.


5 The synthesis of pyrazoles of Formula I can be prepared as shown in Schemes 1 and 2. The boronic acid coupling was accomplished using the procedure of Lam, Y. S. *et al. Tetrahedron Lett.* 39: 2941-2944 (1998).

Scheme 1



10

Scheme 2




Triazoles of Formula I can be prepared as shown in Scheme 3, employing commercially available 4-(1,2,4-triazol-1-yl)phenol (Lancaster Synthesis).

15

- 16 -

Scheme 3



The invention is also directed to ³H and ¹⁴C radiolabeled compounds of Formula *I* and their use as radioligands for their binding site on the sodium channel.

For example, one use of the labeled compounds of the invention is the characterization of specific receptor binding. Another use of the labeled compounds of the invention is an alternative to animal testing for the evaluation of structure-activity relationships. The receptor assay is performed at a fixed concentration of a labeled compound of Formula *I* and at increasing concentrations of a test compound in a competition assay.

Tritiated compounds of Formula *I* can be prepared by introducing tritium into the compound of Formula *I* by, for example, catalytic dehalogenation with tritium. This method includes reacting a suitably halogen-substituted precursor of a compound of Formula *I* with tritium gas in the presence of a suitable catalyst, for example Pd/C, in the presence or absence of a base. Other suitable methods for preparing tritiated compounds can be found in Filer, *Isotopes in the Physical and Biomedical Sciences, Vol. 1, Labeled Compounds (Part A)*, Chapter 6. ¹⁴C-labeled compounds can be prepared by employing starting materials having a ¹⁴C carbon.

The compounds of the present invention can be assessed by electrophysiological assays in dissociated hippocampal neurons for sodium channel blocker activity. These compounds also can be assayed for binding to the neuronal voltage-dependent sodium channel using rat forebrain membranes and [³H]BTX-B.

Sodium channels are large transmembrane proteins that are expressed in various tissues. They are voltage sensitive channels and are responsible for the rapid increase of Na⁺ permeability in response to depolarization associated with the action potential in many excitable cells including muscle, nerve and cardiac cells.

One aspect of the present invention is the discovery of the mechanism of action of the compounds herein described as specific Na^+ channel blockers. Based upon the discovery of this mechanism, these compounds are contemplated to be useful in treating or preventing neuronal loss due to focal or global ischemia, and in treating or preventing neurodegenerative disorders including ALS, anxiety, and epilepsy. They are also expected to be effective in treating, preventing or ameliorating neuropathic pain, surgical pain, chronic pain and tinnitus. The compounds are also expected to be useful as antiarrhythmics, anesthetics and antimanic depressants.

The present invention is directed to compounds of Formula I that are blockers of voltage-sensitive sodium channels. According to the present invention, those compounds having preferred sodium channel blocking properties exhibit an IC_{50} of about 100 μM or less in the electrophysiological assay described herein. Preferably, the compounds of the present invention exhibit an IC_{50} of 10 μM or less. Most preferably, the compounds of the present invention exhibit an IC_{50} of about 1.0 μM or less. Substituted heteroaryl compounds of the present invention may be tested for their Na^+ channel blocking activity by the following electrophysiological and binding assays.

Electrophysiological Assay:

Cell preparation: HEK-293 (NaIIA-B2) cell line stably expressing the rBIIA isoform of Na^+ channels is established in-house. The cells are cultured using standard techniques, as described previously (Verdoorn, T.A., *et al.*, *Neuron* 4:919-928 (1990)). For electrophysiology, cells are plated onto poly-D-lysine pre-coated Cellware 35 mm Petri dishes (BIOCOAT, Becton Dickinson) at a density of $\sim 10^4$ cells/dish on the day of re-seeding from confluent cultures. Our experience has been that cells are suitable for recordings for 2-3 days after plating.

Patch-clamp recordings of voltage-sensitive Na^+ currents: Whole-cell voltage-clamp recordings are made using conventional patch-clamp techniques (Hamill *et al.*, *Pfluegers Arch.* 391:85-100 (1981)) with an Axopatch 200A amplifier (Axon Instruments, Foster City, CA). The recording chamber is continuously

- 18 -

superfused with the external solution (150 mM NaCl, 5.4 mM KCl, 1.8 mM CaCl_2 , 1 mM MgCl_2 , 10 mM HEPES, 10 mM glucose, pH 7.4 adjusted with NaOH, osmolality ~ 320 mmol/kg) at a speed of about 1 mL/min. Recording pipettes were pulled from the thick-walled capillaries (WPI, Sarasota, FL) and fire-polished. The pipette resistances range from 1 to 3 M Ω when the pipettes are filled with internal solution containing (in mM): 130 CsF, 20 NaCl, 2 MgCl_2 , 10 EGTA, 10 HEPES, pH adjusted to 7.4 with CsOH, osmolality ~ 310 mmol/kg. Drugs and intervening wash-outs are applied through a linear array of flow pipes (Drummond Microcaps, 2 μL , 64-mm length). Compounds are dissolved in dimethylsulfoxide (DMSO) to make a 30 mM stock solution, which is subsequently diluted into the external solution to give final concentrations of 0.1-100 μM . At the highest (1 %) concentration, DMSO inhibits the size of Na^+ current only slightly. Currents are recorded at room temperature (22-25 $^{\circ}\text{C}$), filtered at 3 kHz with an active 8-pole Bessel filter (Frequency Devices, Haverhill, MA), digitized at 10-50 μs intervals, and stored using Digidata 1200 analog/digital interface with Pclamp6/Clampex software (Axon Instruments). Series resistance is cancelled typically by $\sim 75\%$ when necessary.

The following voltage pulse protocols are used to assess the potency and kinetics of inhibition of the Na^+ channels by the compounds (Fig. 1).




Figure 1. Voltage pulse protocols. A. IV-curves. C. Steady-state inactivation. B. Repriming kinetics. D. Time course of binding.

Current-voltage relationship (IV-curve), protocol A, is used to report the voltage at which the maximal inward Na^+ current is achieved. This voltage is used throughout

the experiment as testing voltage, V_t . The steady-state inactivation (or, availability) curve, protocol C, is used to get the voltage at which almost complete ($\geq 95\%$) inactivation of Na^+ channels occurs; it serves as voltage for conditioning prepulse, V_c , throughout the experiment. Protocol B reports how fast the channels recover from inactivation at hyperpolarized voltages. This permits us to set up the duration of the hyperpolarization gap which is used in measurement of the kinetics of binding of compounds to inactivated Na^+ channels (protocol D). Channel repriming under control conditions is fast ($\geq 90\%$ recovery during first 5-10 ms). If a drug substantially retards the repriming process, then it becomes possible (protocol D) to accurately measure the kinetics of binding of the inhibitor to inactivated channels as well as the steady-state affinity (k_+ and K_i). To estimate k_+ values, the reduction in peak currents in successive trials with varying pre-pulse duration is plotted as a function of pre-pulse duration and the time constant (τ) measured by mono-exponential fit. A plot of $1/\tau$ as a function of antagonist concentration then allows calculating of the macroscopic binding rates of the antagonists. To determine K_i values the partial inhibition curves measured by fractional responses in steady-state are fitted with the logistic equation:

$$I/I_{\text{control}} = 1/(1 + ([\text{antagonist}]/K_i)^p), \quad \text{Eq. 2}$$

where I_{control} is the maximal Na^+ current in the absence of antagonist, $[\text{antagonist}]$ is the drug concentration, K_i is the concentration of antagonist that produces half maximal inhibition, and p is the slope factor.

In vitro Binding Assay:

The ability of compounds of the present invention to modulate either site 1 or site 2 of the Na^+ channel is determined following the procedures fully described in Yasushi, *J. Biol. Chem.* 261:6149-6152 (1986) and Creveling, *Mol. Pharmacol.* 23:350-358 (1983), respectively. Rat forebrain membranes are used as sources of Na^+ channel proteins. The binding assays are conducted in 130 μM choline chloride at 37

- 20 -

°C for 60-minute incubation with [³H] saxitoxin and [³H] batrachotoxin as radioligands for site 1 and site 2, respectively.

In vivo Pharmacology:

5

The compounds of the present invention may be tested for *in vivo* anticonvulsant activity after i.v., p.o. or i.p. injection using a number of anticonvulsant tests in mice, including the maximum electroshock seizure test (MES). Maximum electroshock seizures are induced in male NSA mice weighing between 15-20 g and male Sprague-Dawley rats weighing between 200-225 g by application of current (50
10 mA, 60 pulses/sec, 0.8 msec pulse width, 1 sec duration, D.C., mice; 99 mA, 125 pulses/sec, 0.8 msec pulse width, 2 sec duration, D.C., rats) using a Ugo Basile ECT device (Model 7801). Mice are restrained by gripping the loose skin on their dorsal surface and saline-coated corneal electrodes are held lightly against the two corneae.
15 Rats are allowed free movement on the bench top and ear-clip electrodes are used. Current is applied and animals are observed for a period of up to 30 seconds for the occurrence of a tonic hindlimb extensor response. A tonic seizure is defined as a hindlimb extension in excess of 90 degrees from the plane of the body. Results are treated in a quantal manner.

20

The compounds may be tested for their antinociceptive activity in the formalin model as described in Hunskaar, S., O. B. Fasmer, and K. Hole, *J. Neurosci. Methods* 14: 69-76 (1985). Male Swiss Webster NIH mice (20-30 g; Harlan, San Diego, CA) are used in all experiments. Food is withdrawn on the day of experiment. Mice are placed in Plexiglass jars for at least 1 hour to accommodate to the environment.
25 Following the accommodation period mice are weighed and given either the compound of interest administered i.p. or p.o., or the appropriate volume of vehicle (10 % Tween-80). Fifteen minutes after the i.p. dosing, and 30 minutes after the p.o. dosing mice are injected with formalin (20 µL of 5% formaldehyde solution in saline) into the dorsal surface of the right hind paw. Mice are transferred to the Plexiglass
30 jars and monitored for the amount of time spent licking or biting the injected paw. Periods of licking and biting are recorded in 5 minute intervals for 1 hour after the

formalin injection. All experiments are done in a blinded manner during the light cycle. The early phase of the formalin response is measured as licking / biting between 0-5 min, and the late phase is measured from 15-50 min. Differences between vehicle and drug treated groups are analyzed by one-way analysis of variance (ANOVA). A P value ≤ 0.05 is considered significant. Having activity in blocking the acute and second phase of formalin-induced paw-licking activity, the compounds are considered to be efficacious for acute and chronic pain.

The compounds may be tested for their potential for the treatment of chronic pain (antiallodynic and antihyperalgesic activities) in the Chung model of peripheral neuropathy. Male Sprague-Dawley rats weighing between 200-225 g are anesthetized with halothane (1-3 % in a mixture of 70 % air and 30 % oxygen) and their body temperature is controlled during anesthesia through use of a homeothermic blanket. A 2-cm dorsal midline incision is then made at the L5 and L6 level and the paravertebral muscle groups are retracted bilaterally. L5 and L6 spinal nerves are then exposed, isolated, and tightly ligated with 6-0 silk suture. A sham operation is performed exposing the contralateral L5 and L6 spinal nerves as a negative control.

Tactile Allodynia: Rats are transferred to an elevated testing cage with a wire mesh floor and allowed to acclimate for five to ten minutes. A series of Semmes-Weinstein monofilaments are applied to the plantar surface of the hindpaw to determine the animal's withdrawal threshold. The first filament used possesses a buckling weight of 9.1 gms (.96 log value) and is applied up to five times to see if it elicits a withdrawal response. If the animal has a withdrawal response then the next lightest filament in the series would be applied up to five times to determine if it could elicit a response. This procedure is repeated with subsequent lesser filaments until there is no response and the lightest filament that elicited a response is recorded. If the animal does not have a withdrawal response from the initial 9.1 gms filament then subsequent filaments of increased weight are applied until a filament elicits a response and this filament is then recorded. For each animal, three measurements are made at every time point to produce an average withdrawal threshold determination. Tests are performed prior to and at 1, 2, 4 and 24 hours post-drug administration. Tactile allodynia and mechanical hyperalgesia tests are conducted concurrently.

- 22 -

Mechanical Hyperalgesia: Rats are transferred to an elevated testing cage with a wire mesh floor and allowed to acclimate for five to ten minutes. A slightly blunted needle is touched to the plantar surface of the hindpaw causing a dimpling of the skin without penetrating the skin. Administration of the needle to control paws typically produces a quick flinching reaction, too short to be timed with a stopwatch and arbitrarily given a withdrawal time of 0.5 sec. The operated side paw of neuropathic animals exhibits an exaggerated withdrawal response to the blunted needle. A maximum withdrawal time of ten seconds is used as a cutoff time. Withdrawal times for both paws of the animals are measured three times at each time point with a five-minute recovery period between applications. The three measures are used to generate an average withdrawal time for each time point. Tactile allodynia and mechanical hyperalgesia tests are conducted concurrently.

The compounds may be tested for their neuroprotective activity after focal and global ischemia produced in rats or gerbils according to the procedures described in Buchan *et al.* (*Stroke*, Suppl. 148-152 (1993)) and Sheardown *et al.* (*Eur. J. Pharmacol.* 236:347-353 (1993)) and Graham *et al.* (*J. Pharmacol. Exp. Therap.* 276:1-4 (1996)).

The compounds may be tested for their neuroprotective activity after traumatic spinal cord injury according to the procedures described in Wrathall *et al.* (*Exp. Neurology* 137:119-126 (1996)) and Iwasaki *et al.* (*J. Neuro Sci.* 134:21-25 (1995)).

Compositions within the scope of this invention include all compositions wherein the compounds of the present invention are contained in an amount that is effective to achieve its intended purpose. While individual needs vary, determination of optimal ranges of effective amounts of each component is within the skill of the art. Typically, the compounds may be administered to mammals, e.g. humans, orally at a dose of 0.0025 to 50 mg/kg, or an equivalent amount of the pharmaceutically acceptable salt thereof, per day of the body weight of the mammal being treated for epilepsy, neurodegenerative diseases, anesthetic, arrhythmia, manic depression, and pain. For intramuscular injection, the dose is generally about one-half of the oral dose.

In the method of treatment or prevention of neuronal loss in global and focal ischemia, brain and spinal cord trauma, hypoxia, hypoglycemia, status epilepsy and surgery, the compound can be administrated by intravenous injection at a dose of about 0.025 to about 10 mg/kg.

5 The unit oral dose may comprise from about 0.01 to about 50 mg, preferably about 0.1 to about 10 mg of the compound. The unit dose may be administered one or more times daily as one or more tablets each containing from about 0.1 to about 10, conveniently about 0.25 to 50 mg of the compound or its solvates.

10 In addition to administering the compound as a raw chemical, the compounds of the invention may be administered as part of a pharmaceutical preparation containing suitable pharmaceutically acceptable carriers comprising excipients and auxiliaries which facilitate processing of the compounds into preparations which can be used pharmaceutically. Preferably, the preparations, particularly those preparations
15 which can be administered orally and which can be used for the preferred type of administration, such as tablets, dragees, and capsules, and also preparations which can be administered rectally, such as suppositories, as well as suitable solutions for administration by injection or orally, contain from about 0.01 to 99 percent, preferably from about 0.25 to 75 percent of active compound(s), together with the excipient.

20 Also included within the scope of the present invention are the non-toxic pharmaceutically acceptable salts of the compounds of the present invention. Acid addition salts are formed by mixing a solution of the particular heteroaryl compound of the present invention with a solution of a pharmaceutically acceptable non-toxic acid such as hydrochloric acid, fumaric acid, maleic acid, succinic acid, acetic acid, citric acid, tartaric acid, carbonic acid, phosphoric acid, oxalic acid, dichloroacetic
25 acid, and the like. Basic salts are formed by mixing a solution of the heteroaryl compound of the present invention with a solution of a pharmaceutically acceptable non-toxic base such as sodium hydroxide, potassium hydroxide, choline hydroxide, sodium carbonate and the like.

30 The pharmaceutical compositions of the invention may be administered to any animal that may experience the beneficial effects of the compounds of the invention.

- 24 -

Foremost among such animals are mammals, e.g., humans, although the invention is not intended to be so limited.

5 The pharmaceutical compositions of the present invention may be administered by any means that achieve their intended purpose. For example, administration may be by parenteral, subcutaneous, intravenous, intramuscular, intraperitoneal, transdermal, or buccal routes. Alternatively, or concurrently, administration may be by the oral route. The dosage administered will be dependent upon the age, health, and weight of the recipient, kind of concurrent treatment, if any, frequency of treatment, and the nature of the effect desired.

10 The pharmaceutical preparations of the present invention are manufactured in a manner which is itself known, for example, by means of conventional mixing, granulating, dragee-making, dissolving, or lyophilizing processes. Thus, pharmaceutical preparations for oral use can be obtained by combining the active compounds with solid excipients, optionally grinding the resulting mixture and
15 processing the mixture of granules, after adding suitable auxiliaries, if desired or necessary, to obtain tablets or dragee cores.

Suitable excipients are, in particular, fillers such as saccharides, for example lactose or sucrose, mannitol or sorbitol, cellulose preparations and/or calcium phosphates, for example tricalcium phosphate or calcium hydrogen phosphate, as well
20 as binders such as starch paste, using, for example, maize starch, wheat starch, rice starch, potato starch, gelatin, tragacanth, methyl cellulose, hydroxypropylmethylcellulose, sodium carboxymethylcellulose, and/or polyvinyl pyrrolidone. If desired, disintegrating agents may be added such as the above-mentioned starches and also carboxymethyl-starch, cross-linked polyvinyl pyrrolidone, agar, or alginic
25 acid or a salt thereof, such as sodium alginate. Auxiliaries are, above all, flow-regulating agents and lubricants, for example, silica, talc, stearic acid or salts thereof, such as magnesium stearate or calcium stearate, and/or polyethylene glycol. Dragee cores are provided with suitable coatings which, if desired, are resistant to gastric juices. For this purpose, concentrated saccharide solutions may be used, which may
30 optionally contain gum arabic, talc, polyvinyl pyrrolidone, polyethylene glycol and/or titanium dioxide, lacquer solutions and suitable organic solvents or solvent mixtures.

In order to produce coatings resistant to gastric juices, solutions of suitable cellulose preparations such as acetylcellulose phthalate or hydroxypropymethyl-cellulose phthalate, are used. Dye stuffs or pigments may be added to the tablets or dragee coatings, for example, for identification or in order to characterize combinations of active compound doses.

Other pharmaceutical preparations which can be used orally include push-fit capsules made of gelatin, as well as soft, sealed capsules made of gelatin and a plasticizer such as glycerol or sorbitol. The push-fit capsules can contain the active compounds in the form of granules which may be mixed with fillers such as lactose, binders such as starches, and/or lubricants such as talc or magnesium stearate and, optionally, stabilizers. In soft capsules, the active compounds are preferably dissolved or suspended in suitable liquids, such as fatty oils, or liquid paraffin. In addition, stabilizers may be added.

Possible pharmaceutical preparations, which can be used rectally, include, for example, suppositories, which consist of a combination of one or more of the active compounds with a suppository base. Suitable suppository bases are, for example, natural or synthetic triglycerides, or paraffin hydrocarbons. In addition, it is also possible to use gelatin rectal capsules which consist of a combination of the active compounds with a base. Possible base materials include, for example, liquid triglycerides, polyethylene glycols, or paraffin hydrocarbons.

Suitable formulations for parenteral administration include aqueous solutions of the active compounds in water-soluble form, for example, water-soluble salts and alkaline solutions. In addition, suspensions of the active compounds as appropriate oily injection suspensions may be administered. Suitable lipophilic solvents or vehicles include fatty oils, for example, sesame oil, or synthetic fatty acid esters, for example, ethyl oleate or triglycerides or polyethylene glycol-400 (the compounds are soluble in PEG-400). Aqueous injection suspensions may contain substances which increase the viscosity of the suspension, and include, for example, sodium carboxymethyl cellulose, sorbitol, and/or dextran. Optionally, the suspension may also contain stabilizers.

- 26 -

The following examples are illustrative, but not limiting, of the method and compositions of the present invention. Other suitable modifications and adaptations of the variety of conditions and parameters normally encountered in clinical therapy and which are obvious to those skilled in the art are within the spirit and scope of the invention.

1-[4-(4-Fluorophenoxy)phenyl]-3-methylpyrazole

a). **4-(4-Fluorophenoxy)phenylhydrazine hydrochloride.** A suspension of finely powdered 4-fluoro-4'-aminodiphenyl ether (2.00 g, 9.84 mmol) in 10 mL of water was cooled in an ice-water bath and 19.4 mL of conc. HCl was added dropwise via addition funnel. The resulting mixture was cooled to -5 °C in an acetone-ice bath and a solution of sodium nitrite (crystalline; 0.714 g, 10.3 mmol) in 8 mL of cold water was added dropwise to the reaction at such a rate that the temperature remained between and -5 and 0 °C. A solution of SnCl₂·2H₂O (6.66 g, 29.5 mmol) in 20 mL of conc. HCl at -20 °C was treated with the reaction mixture added in portions, maintaining the temperature below -10 °C. A grey ppt. formed immediately and the resulting mixture was stirred at -20 °C for 90 min. The solid was isolated by filtration and washed with cold EtOH (2 x 10 mL). The crude hydrazine, 2.36 g, was carried on without purification.

b). **1-[4-(4-Fluorophenoxy)phenyl]-3-methylpyrazole.** A suspension of the hydrazine (500 mg, 2.05 mmol) in 5.5 mL of 1:1 EtOH/water was treated with 300 µL (299 mg, 2.03 mmol) of 90% acetylacetaldehyde dimethylacetal and the resulting mixture was warmed with a heat gun for 2 min. The reaction was allowed to cool and extracted with hexane (4 x 10 mL). The pooled organic layers were washed with brine, dried (Na₂SO₄) and concentrated. The residue was subjected to column chromatography (silica gel, 10% EtOAc/hexane) affording 134 mg (24%) of the title compound as a white solid, mp 80-81 °C. ¹H NMR (CDCl₃): δ 7.74 (s, 1H), 7.58 (d, 2H, J = 8.4 Hz), 7.06-6.95 (m, 6H), 6.23 (s, 1H), 2.37 (s, 3H).

1-(4-Phenoxyphenyl)-1H-pyrazole-3-carboxamide and 1-(4-phenoxyphenyl)-1H-pyrazole-5-carboxamide

a). 3-Ethoxycarbonyl-1-(4-phenoxyphenyl)-1H-pyrazole and 5-ethoxycarbonyl-

1-(4-phenoxyphenyl)-1H-pyrazole. To a suspension of 4-phenoxyphenylboronic acid (1.70 g, 7.85 mmol), ethyl 3-pyrazolecarboxylate (0.55 g, 3.92 mmol), copper(II)acetate (1.1 g, 5.89 mmol) and 4Å molecular sieves (powdered and heated at 200 °C for 2h prior to use) in 30 mL of anhydrous THF was added 0.6 mL of pyridine. The reaction was stirred open to air at room temperature for 2 days and then filtered and the filtrate was concentrated to dryness. The crude product was purified by flash chromatography, eluting with 15% EtOAc/hexane, affording 5-ethoxycarbonyl-1-(4-phenoxyphenyl)-1H-pyrazole (R_f = 0.6, 55 mg, 4.6%) and 3-ethoxycarbonyl-1-(4-phenoxyphenyl)-1H-pyrazole (R_f = 0.5, 125 mg, 10.3%).

b). 1-(4-Phenoxyphenyl)-1H-pyrazole-3-carboxamide. A solution of 3-ethoxycarbonyl-1-(4-phenoxyphenyl)-1H-pyrazole (120 mg, 0.39 mmol) in 5 mL of a 2N solution of ammonia in MeOH was stirred at rt for 4 d. TLC showed incomplete reaction and the solution was transferred to a sealed tube and heated at 70 °C overnight. The reaction was concentrated and purified by preparative TLC, eluting with 50% EtOAc/hexane, to yield 1-(4-phenoxyphenyl)-1H-pyrazole-3-carboxamide (R_f = 0.26, 56 mg, 52%), mp 165-167 °C. ^1H NMR (300 MHz, DMSO- d_6) δ 8.50 (d, J = 2.7 Hz, 1H, pyrazole), 7.92 (d, J = 9.0 Hz, 2H, Phenyl), 7.71 (br, 1H, NH_2), 7.43 (m, 2H, Phenoxy), 7.39 (br, 1H, NH_2), 7.18 (m, 1H, Phenoxy), 7.17 (d, J = 9.0 Hz, 2H, Phenyl), 7.07 (m, 2H, Phenoxy), 6.87 (d, J = 2.7 Hz, 1H, pyrazole).

Starting with 5-ethoxycarbonyl-1-(4-phenoxyphenyl)-1H-pyrazole (50 mg, 0.16 mmol), the method described above gave 1-(4-phenoxyphenyl)-1H-pyrazole-5-carboxamide (25 mg, 55%), mp 142-144 °C. ^1H NMR (300 MHz, DMSO- d_6) δ 8.03 (br s, 1H, NH_2), 7.70 (d, J = 1.8 Hz, 1H, pyrazole), 7.54 (br s, 1H, NH_2), 7.42 (m, 2H, Phenoxy), 7.39 (d, J = 9.0 Hz, 2H, Phenyl), 7.20 (m, 1H, Phenoxy), 7.08 (m, 2H, Phenoxy), 7.06 (d, J = 9.0 Hz, 2H, Phenyl), 6.90 (d, J = 1.8 Hz, 1H, pyrazole).

- 28 -

1-[4-(4-Nitrophenoxy)phenyl]-1H-[1,2,4]triazole

A mixture of 1-fluoro-4-nitrobenzene (0.17 mL, 1.6 mmol), 4-{[1,2,4]triazol-1-yl}phenol (0.26 g, 1.58 mmol), and potassium carbonate (1.69 g, 12.2 mmol) in DMF was refluxed overnight. The reaction was cooled to room temperature, then partitioned between water and ethyl acetate. The aqueous layer was extracted with ethyl acetate. The combined organic layers were washed with an aqueous sodium hydroxide solution (2N), water (2 times), dried over sodium sulfate, filtered, and evaporated under reduced pressure to give a yellow solid. Purification by column chromatography (silica gel; 1:1 hexane/ethyl acetate) and recrystallization from chloroform/hexane afforded 165 mg (37%) of the title compound as a yellow solid, mp 131-132 °C. ¹H-NMR (CDCl₃): δ 8.55 (s, 1H), 8.25 (d, J = 9 Hz, 2H), 8.12 (s, 1H), 7.75 (d, J = 9 Hz, 2H), 7.24 (d, J = 9 Hz, 2H), 7.08 (d, J = 9 Hz, 2H).

Anticonvulsant Activity of Compounds of the Invention

The ability of compounds of the present invention to block maximal electroshock-induced seizures (MES) is determined as described earlier.

A compound of the present invention is administered p.o. to mice 30 minutes before the test procedure. The compound exhibits protection against MES with an ED₅₀ (the dose provided protection of 50% of animals) of preferably below 10 mg/kg.

Activity of 1-(4-phenoxyphenyl)-1H-pyrazole-3-carboxamide as anticonvulsant, MES i.v. ED₅₀ in mouse 0.7 mg/kg. 1-[4-(4-Nitrophenoxy)phenyl]-1H-[1,2,4]triazole, MES p.o. ED₅₀ 6.6 mg/kg. 1-(4-phenoxyphenyl)-1H-pyrazole-5-carboxamide, MES i.v. ED₅₀ 10.0 mg/kg.

Activity of Compound of the Invention as Sodium Channel Blocker

5 Compounds of the invention are tested in the electrophysiological and binding assays described above and produce dose-dependent inhibition of voltage-gated sodium currents recorded in HEK-293 cells stably expressing the rBIIA isoform of Na⁺ channels. The blocking effect of preferred compounds on Na⁺ currents is highly sensitive to the holding voltage, indicating that the compounds bind to voltage-
10 sensitive Na⁺ channels in their inactivated states and have weak potency towards Na⁺ channels in their resting states (Ragsdale *et al.*, *Mol. Pharmacol.* 40:756-765 (1991); Kuo and Bean, *Mol. Pharmacol.* 46:716-725 (1994)). The apparent antagonist dissociation constant (K_d) of preferred compounds for inactivated sodium channels is less than 400 nM. 1-(4-Phenoxyphenyl)-1H-pyrazole-3-carboxamide was tested in
15 rBIIA isoform of sodium channel and had K_i 0.35 μ M.

 Having now fully described this invention, it will be understood by those of ordinary skill in the art that the same can be performed within a wide and equivalent
20 range of conditions, formulations and other parameters without affecting the scope of the invention or any embodiment thereof. All patents and publications cited herein are fully incorporated by reference herein in their entirety.

What Is Claimed Is:

1. A compound having the Formula I:




5

or a pharmaceutically acceptable salt, prodrug or solvate thereof, wherein

X is one of O, S, NR₉, CH₂, NR₉C(O), or C(O)NR₉, where R₉ is hydrogen or C₁-C₁₀ alkyl;

Het is a heteroaryl selected from the group consisting of

10



R₁ is selected from the group consisting of hydrogen, optionally substituted alkyl, optionally substituted heteroaryl, C(O)R₁₀, CH₂C(O)R₁₀, S(O)R₁₀, and SO₂R₁₀;

15

R₂ and R₃ are independently selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, aryl, cyano, aminoalkyl, hydroxyalkyl, alkoxyalkyl, alkylthio, alkylsulfinyl, alkylsulfonyl, carboxyalkyl, alkylamino, dialkylamino, aminocarbonyl, alkylaminocarbonyl, arylaminocarbonyl, aralkylaminocarbonyl, alkylcarbonylamino, arylcarbonylamino, aralkylcarbonylamino, alkylcarbonyl, aminosulfonyl, alkylaminosulfonyl, and alkylsulfonyl;

20

R₅, R₆, R₇, and R₈ are independently selected from the group consisting of hydrogen, halo, haloalkyl, alkyl, alkenyl, alkynyl, hydroxyalkyl, aminoalkyl, carboxyalkyl, alkoxyalkyl, nitro, amino, ureido, cyano, acylamino, amide, hydroxy, thiol, acyloxy, azido, alkoxy, carboxy, carbonylamido and alkylthiol;

R_{10} is selected from the group consisting of amino, alkyl, alkenyl, alkynyl, OR_{11} , alkylamino, dialkylamino, alkenylamino, dialkylaminoalkenyl, cycloalkyl, heterocycle, heteroaryl, aryl, aralkyl, arylalkenyl, arylalkynyl, and cycloalkylalkylamino;

5 R_{11} is selected from the group consisting of hydrogen, optionally substituted alkyl, and an alkalimetal; and

provided that:

- 1) when Het is (ii), and X is O, then R_{10} is not alkyl, aralkyl, aryl or OR_{11} ;
- 2) when Het is (i) or (ii), then X is not NR_9 ;
- 10 3) when Het is (iii), then X is not CH_2 ; and
- 4) when Het is (iii), and X is O, then R_{10} is not OR_{11} .

2. A compound of claim 1, wherein R_1 is selected from the group
15 consisting of an alkyl optionally substituted by halogen, hydroxy, carbamoyloxy, C_{1-6} acyl, C_{1-6} alkylsulfonylamino, aryl, preferably phenyl, or aminocarbonyl; $C(O)R_{10}$; $CH_2C(O)R_{10}$; or SO_2R_{10} , wherein R_{10} is selected from the group consisting of C_{1-6} alkyl, C_{2-6} alkenyl, OR_{11} , amino, C_{1-6} alkylamino, $di(C_{1-6})$ alkylamino, C_{2-6} alkenylamino, heterocycle and mono- and di- (C_{1-6}) alkylaminoalkenyl, and R_{11} is
20 selected from the group consisting of hydrogen, optionally substituted alkyl, and an alkalimetal.

3. A compound of claim 2, wherein R_{10} is selected from the group
consisting of C_{1-6} alkyl, C_{2-6} alkenyl, OR_{10} , amino, C_{1-6} alkylamino, $di(C_{1-6})$ alkylamino,
25 C_{2-6} alkenylamino, $di(C_{1-6})$ alkylamino(C_{2-6})alkenyl, N-morpholinyl, N-pyrrolidinyl, and N-piperazinyl.

4. A compound of claim 3, wherein R_2 and R_3 are independently selected
from the group consisting of hydrogen, C_1-C_6 alkyl, C_2-C_6 alkenyl, C_2-C_6 alkynyl,
30 amino(C_1-C_6)alkyl, amino, C_1-C_6 alkylthio, cyano, C_1-C_6 alkylsulfinyl, hydroxy(C_1-C_6)alkyl, C_1-C_6 alkoxy, aminocarbonyl, C_1-C_6 alkylaminocarbonyl, C_6-C_{10} arylaminocarbonyl, C_6-C_{10} aryl(C_1-C_6)alkylaminocarbonyl, C_1-C_6 alkylcarbonylamino, C_6-C_{10} arylcarbonylamino, and C_6-C_{10} aryl(C_1-C_6)alkylcarbonylamino.

- 32 -

5. A compound of claim 3, wherein R_2 and R_3 are independently selected from the group consisting of hydrogen, C_1 - C_6 alkyl, C_1 - C_6 alkoxy, amino(C_1 - C_6)alkyl, C_1 - C_6 alkylthio and aminocarbonyl.

5

6. A compound of claim 1, wherein R_5 , R_6 , R_7 , and R_8 are independently selected from the group consisting of hydrogen, halo, halo(C_1 - C_6)alkyl, C_1 - C_6 alkyl, hydroxy(C_1 - C_6)alkyl, amino(C_1 - C_6)alkyl, carboxy(C_1 - C_6)alkyl, alkoxy(C_1 - C_6)alkyl, nitro, amino, C_1 - C_6 acylamino, amide, hydroxy, thiol, C_1 - C_6 acyloxy, C_1 - C_6 alkoxy, carboxy, carbonylamido and C_1 - C_6 alkylthiol.

10

7. A compound of claim 1, wherein R_1 or R_2 is $C(O)R_{10}$ or SO_2R_{10} .

8. A compound of claim 7, wherein where R_{10} is amino or C_{1-6} alkyl.

15

9. A compound of claim 8, wherein X is O or S.

10. A compound of claim 9, wherein:

R_5 and R_6 are each hydrogen;

20

R_3 and R_4 are both H; and

R_7 and R_8 are selected from the group consisting of hydrogen, halo, halo(C_1 - C_6)alkyl, C_1 - C_6 alkyl, hydroxy(C_1 - C_6)alkyl, amino(C_1 - C_6)alkyl, carboxy(C_1 - C_6)alkyl, alkoxy(C_1 - C_6)alkyl, nitro, amino, C_1 - C_6 acylamino, amide, hydroxy, thiol, C_1 - C_6 acyloxy, C_1 - C_6 alkoxy, carboxy, carbonylamido and C_1 - C_6 alkylthiol.

25

11. A compound of claim 10, wherein Het is (i).

12. A compound of claim 10, wherein Het is (ii).

30

13. A compound of claim 10, wherein Het is (iii).

14. A compound of claim 10, wherein Het is (iv).

15. A compound of claim 1, wherein:

Het is (i), (ii), (iii) or (vi);

R_1 is $C(O)R_{10}$, $CH_2C(O)R_{10}$, or SO_2R_{10} ;

X is O or S;


R_{10} is amino, optionally substituted C_1 - C_6 alkyl, or a heterocycle selected from the group consisting of N-morpholinyl, N-pyrrolidinyl and N-piperazinyl;

R_2 , and R_3 are independently hydrogen, C_1 - C_6 alkyl, C_1 - C_6 alkylthio or C_1 - C_6 alkylsulfinyl,

R_5 and R_6 are as defined above and are preferably hydrogen, and

R_7 and R_8 are independently selected from the group consisting of hydrogen, halo, halo(C_1 - C_6)alkyl, C_1 - C_6 alkyl, hydroxy(C_1 - C_6)alkyl, amino(C_1 - C_6)alkyl, carboxy(C_1 - C_6)alkyl, alkoxy(C_1 - C_6)alkyl, nitro, amino, C_1 - C_6 acylamino, amide, hydroxy, thiol, C_1 - C_6 acyloxy, C_1 - C_6 alkoxy, carboxy, carbonylamido and C_1 - C_6 alkylthiol.

16. A compound of Formula I:




I

or a pharmaceutically acceptable salt, prodrug or solvate thereof, wherein

X is O or S;

Het is a heteroaryl selected from the group consisting of

- 34 -



R_1 is $C(O)R_{10}$, $CH_2C(O)R_{10}$, or SO_2R_{10} wherein R_{10} is amino, alkyl, N-morpholinyl, N-pyrrolidinyl or N-piperazinyl, all of which can be optionally substituted;

5 R_2 and R_3 are independently hydrogen, C_1 - C_6 alkyl, C_1 - C_6 alkylthio or C_1 - C_6 alkylsulfinyl;


R_5 , R_6 , R_7 and R_8 are independently selected from the group consisting of hydrogen, halo, halo(C_1 - C_6)alkyl, C_1 - C_6 alkyl, hydroxy(C_1 - C_6)alkyl, amino(C_1 - C_6)alkyl, carboxy(C_1 - C_6)alkyl, alkoxy(C_1 - C_6)alkyl, nitro, amino, C_1 - C_6 acylamino, amide, hydroxy, thiol, C_1 - C_6 acyloxy, C_1 - C_6 alkoxy, carboxy, carbonylamido and C_1 - C_6 alkylthiol;

provided that:

- 1) when Het is (ii), and X is O, then R_{10} is not alkyl, aralkyl, aryl or OR_{11} ; and
- 2) when Het is (iii), and X is O, then R_{10} is not OR_{11} .

17. A pharmaceutical composition, comprising the compound of any one of claims 1-16 and a pharmaceutically acceptable carrier or diluent.


20 18. A method of treating a disorder responsive to the blockade of sodium channels in a mammal suffering therefrom, comprising administering to a mammal in need of such treatment an effective amount of a compound of Formula I:

**I**

25 or a pharmaceutically acceptable salt, prodrug or solvate thereof, wherein

X is one of O, S, NR_9 , CH_2 , $\text{NR}_9\text{C(O)}$, or C(O)NR_9 , where R_9 is hydrogen or $\text{C}_1\text{-C}_{10}$ alkyl;

Het is a heteroaryl selected from the group consisting of



R_1 is selected from the group consisting of hydrogen, optionally substituted alkyl, optionally substituted heteroaryl, C(O)R_{10} , $\text{CH}_2\text{C(O)R}_{10}$, S(O)R_{10} , and SO_2R_{10} ;

10 R_2 and R_3 are independently selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, aryl, cyano, aminoalkyl, hydroxyalkyl, alkoxyalkyl, alkylthio, alkylsulfinyl, alkylsulfonyl, carboxyalkyl, alkylamino, dialkylamino, aminocarbonyl, alkylaminocarbonyl, arylaminocarbonyl, aralkylaminocarbonyl, alkylcarbonylamino, arylcarbonylamino, aralkylcarbonylamino, alkylcarbonyl, aminosulfonyl, alkylaminosulfonyl, and alkylsulfonyl;

15 R_5 , R_6 , R_7 , and R_8 are independently selected from the group consisting of hydrogen, halo, haloalkyl, alkyl, alkenyl, alkynyl, hydroxyalkyl, aminoalkyl, carboxyalkyl, alkoxyalkyl, nitro, amino, ureido, cyano, acylamino, amide, hydroxy, thiol, acyloxy, azido, alkoxy, carboxy, carbonylamido and alkylthiol;

20 R_{10} is selected from the group consisting of amino, alkyl, alkenyl, alkynyl, OR_{11} , alkylamino, dialkylamino, alkenylamino, dialkylaminoalkenyl, cycloalkyl, heterocycle, heteroaryl, aryl, aralkyl, arylalkenyl, arylalkynyl, and cycloalkylalkylamino;

R_{11} is selected from the group consisting of hydrogen, optionally substituted alkyl, and an alkalimetal.

25

19. A method for treating, preventing or ameliorating neuronal loss following global and focal ischemia; treating, preventing or ameliorating neurodegenerative conditions; treating, preventing or ameliorating pain or tinnitus;

- 36 -

treating, preventing or ameliorating manic depression; providing local anesthesia; or treating arrhythmias, or treating convulsions, comprising administering to a mammal in need of such treatment an effective amount of a compound of Formula I:




5

or a pharmaceutically acceptable salt, prodrug or solvate thereof, wherein X is one of O, S, NR₉, CH₂, NR₉C(O), or C(O)NR₉, where R₉ is hydrogen or C₁-C₁₀ alkyl;

Het is a heteroaryl selected from the group consisting of

10



R₁ is selected from the group consisting of hydrogen, optionally substituted alkyl, optionally substituted heteroaryl, C(O)R₁₀, CH₂C(O)R₁₀, S(O)R₁₀, and SO₂R₁₀;

15

R₂ and R₃ are independently selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, aryl, cyano, aminoalkyl, hydroxyalkyl, alkoxyalkyl, alkylthio, alkylsulfinyl, alkylsulfonyl, carboxyalkyl, alkylamino, dialkylamino, aminocarbonyl, alkylaminocarbonyl, arylaminocarbonyl, aralkylaminocarbonyl, alkylcarbonylamino, arylcarbonylamino, aralkylcarbonylamino, alkylcarbonyl, aminosulfonyl, alkylaminosulfonyl, and alkylsulfonyl;

20

R₅, R₆, R₇, and R₈ are independently selected from the group consisting of hydrogen, halo, haloalkyl, alkyl, alkenyl, alkynyl, hydroxyalkyl, aminoalkyl, carboxyalkyl, alkoxyalkyl, nitro, amino, ureido, cyano, acylamino, amide, hydroxy, thiol, acyloxy, azido, alkoxy, carboxy, carbonylamido and alkylthiol;

- 37 -


R_{10} is selected from the group consisting of amino, alkyl, alkenyl, alkynyl, OR_{11} , alkylamino, dialkylamino, alkenylamino, dialkylaminoalkenyl, cycloalkyl, heterocycle, heteroaryl, aryl, aralkyl, arylalkenyl, arylalkynyl, and cycloalkylalkylamino;

5 R_{11} is selected from the group consisting of hydrogen, optionally substituted alkyl, and an alkalimetal.

20. The method of claim 19, wherein the method is for treating, preventing or ameliorating pain and said pain is one of neuropathic pain, surgical pain or
10 chronic pain.

21. A method of alleviating or preventing seizure activity in an animal subject, comprising administering to a mammal in need of such treatment an effective amount of a compound of any of claims 1-16.





22. A compound of Formula I:



or a pharmaceutically acceptable salt, prodrug or solvate thereof, wherein

20 X is O or S, preferably O;

Het is a heteroaryl selected from the group consisting of



- 38 -

R_1 is $C(O)R_{10}$, wherein R_{10} is amino, N-morpholinyl, N-pyrrolidinyl or N-piperazinyl, all of which can be optionally substituted

R_2 and R_3 are independently hydrogen, C_1 - C_6 alkyl, C_1 - C_6 alkylthio or C_1 - C_6 alkylsulfinyl;

- 5 R_5 , R_6 , R_7 and R_8 are independently selected from the group consisting of hydrogen, halo, halo(C_1 - C_6)alkyl, C_1 - C_6 alkyl, hydroxy(C_1 - C_6)alkyl, amino(C_1 - C_6)alkyl, carboxy(C_1 - C_6)alkyl, alkoxy(C_1 - C_6)alkyl, nitro, amino, C_1 - C_6 acylamino, amide, hydroxy, thiol, C_1 - C_6 acyloxy, C_1 - C_6 alkoxy, carboxy, carbonylamido and C_1 - C_6 alkylthiol.

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
4 October 2001 (04.10.2001)

PCT

(10) International Publication Number
WO 01/72714 A3

(51) International Patent Classification⁷: C07D 231/12, 249/08, A61K 31/4196, 31/415, A61P 23/00, 25/08

(21) International Application Number: PCT/US01/08972

(22) International Filing Date: 22 March 2001 (22.03.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/191,757 24 March 2000 (24.03.2000) US

(71) Applicant (for all designated States except US): COCEN-SYS, INC. [US/US]; 213 Technology Drive, Irvine, CA 92618 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): HOGENKAMP, Derk [US/US]; 4510 Park Drive, Carlsbad, CA 92008 (US). NGUYEN, Phong [US/US]; 1048 Wingfoot Street, Placentia, CA 92870 (US). YANG, Ji [CN/US]; 2010 Quail Ridge Drive, Plainsboro, NJ 08536 (US).

(74) Agents: COVERT, John, M. et al.; Sterne, Kessler, Goldstein & Fox P.L.L.C., 1100 New York Avenue, N.W., Suite 600, Washington, DC 20005-3934 (US).

(81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).


Published:

-- with international search report

(88) Date of publication of the international search report:
30 May 2002

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: ARYL SUBSTITUTED PYRAZOLES, TRIAZOLES AND TETRAZOLES AS SODIUM CHANNEL BLOCKER



(57) Abstract: This invention relates to compounds having the Formula (I) or a pharmaceutically acceptable salt, prodrug or solvate thereof, wherein Het and R₅ - R₈ are set in the specification. The invention also is directed to the use of compounds of Formula (I) for the treatment of neuronal damage following global and focal ischemia, for the treatment or prevention of neurodegenerative conditions such as amyotrophic lateral sclerosis (ALS), and for the treatment, prevention or amelioration of both acute or chronic pain, as antitinnitus agents, as anticonvulsants, and as antimanic depressants, as local anesthetics,

as antiarrhythmics and for the treatment or prevention of diabetic neuropathy.

WO 01/72714 A3

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 01/08972

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07D231/12 C07D249/08 A61K31/4196 A61K31/415 A61P23/00
A61P25/08

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D A61K A61P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

CHEM ABS Data, EPO-Internal

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	DATABASE CA 'Online! CHEMICAL ABSTRACTS SERVICE, COLUMBUS, OHIO, US; POLEVOI, L. G. ET AL: "Relation of aminopyrazole structures to their neurotropic activity" retrieved from STN Database accession no. 69:1557 XP002175833 compound with RN=20737-97-7 & IZV. TIMIRYAEV SEL'SKOKHOZ. AKAD. (1968), (1), 192-207 ,	1, 19
X	DE 19 36 760 A (GEIGY) 22 January 1970 (1970-01-22) page 18, line 20 - line 21; claim 1 -/--	1



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance: the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance: the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *G* document member of the same patent family

Date of the actual completion of the international search

27 August 2001

Date of mailing of the international search report

24/09/2001

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

De Jong, B

INTERNATIONAL SEARCH REPORT

In ternational Application No

PLI/US 01/08972

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 96 11911 A (OKUMURA YOSHIYUKI ;PFIZER PHARMA (JP); MANO TAKASHI (JP); NAKAO KA) 25 April 1996 (1996-04-25) claims; example 36 ---	1
P,X	EP 1 081 146 A (PFIZER PROD INC) 7 March 2001 (2001-03-07) page 18, line 24 - line 48; example 10 -----	1,6

FURTHER INFORMATION CONTINUED FROM PCT/SA/ 210

Continuation of Box I.2

Claims Nos.: 1-9,17-21 (all in part)

Present claims 1-9,17-21 relate to an extremely large number of possible compounds. Support within the meaning of Article 6 PCT and/or disclosure within the meaning of Article 5 PCT is to be found, however, for only a very small proportion of the compounds claimed. In the present case, the claims so lack support, and the application so lacks disclosure, that a meaningful search over the whole of the claimed scope is impossible. Consequently, the search has been carried out for those parts of the claims which appear to be supported and disclosed, namely those parts relating to the compounds according to claims 10-16 and 22.

The applicant's attention is drawn to the fact that claims, or parts of claims, relating to inventions in respect of which no international search report has been established need not be the subject of an international preliminary examination (Rule 66.1(e) PCT). The applicant is advised that the EPO policy when acting as an International Preliminary Examining Authority is normally not to carry out a preliminary examination on matter which has not been searched. This is the case irrespective of whether or not the claims are amended following receipt of the search report or during any Chapter II procedure.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 01/08972

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
DE 1936760 A	22-01-1970	CH 498902 A BE 736302 A ES 369617 A FR 2013277 A GB 1277781 A	15-11-1970 19-01-1970 01-06-1971 27-03-1970 14-06-1972
WO 9611911 A	25-04-1996	AT 198474 T AU 681520 B AU 2417495 A BG 61768 B BG 100070 A BR 9504442 A CA 2202057 A CZ 9701177 A DE 69519804 D DE 69519804 T DK 787127 T EP 0787127 A ES 2153033 T FI 971632 A IL 115581 A JP 3032579 B JP 11507322 T KR 239930 B LV 11468 A LV 11468 B NO 971773 A NZ 285163 A PL 319727 A RU 2136666 C SI 9500392 A SK 128795 A TR 960320 A TW 411339 B US 5883106 A US 6063928 A ZA 9508748 A	15-01-2001 28-08-1997 06-05-1996 29-05-1998 31-05-1996 20-05-1997 25-04-1996 17-06-1998 08-02-2001 19-04-2001 05-02-2001 06-08-1997 16-02-2001 17-04-1997 28-10-1999 17-04-2000 29-06-1999 02-03-2000 20-08-1996 20-12-1996 17-06-1997 26-08-1998 18-08-1997 10-09-1999 31-08-1996 03-07-1996 21-06-1996 11-11-2000 16-03-1999 16-05-2000 17-04-1997
EP 1081146 A	07-03-2001	AU 5360700 A CN 1291609 A JP 2001106687 A PL 342284 A	08-03-2001 18-04-2001 17-04-2001 12-03-2001